

A Note on Singularities and Polynomial Zeros

Gordon Chalmers

e-mail: gordon@quartz.shango.com

Abstract

The analysis of solutions to algebraic equations is further simplified. A couple of functions and their analytic continuation or root findings are required.

A simplified version of the solutions to algebraic equations is presented. In previous work, a generating function to sets of these equations is derived via the geodesic flow on potentially singular manifolds [1]. The general solution to these geodesic equations dictates solutions to polynomial equations not only in the integer and rational fields, but also in irrational and transcendental ones. The work [1] is not reviewed here. There is formalism related to the work in [2] and [3] both in number theory and an implementation in scalar field theory.

The solution set to the elliptic equation in

$$y^2 = x^3 + ax + b , \quad (1)$$

and in general the hyperelliptic equations, as well as systems of coupled polynomial equations, is of interest for many reasons. The counting of these solutions can be done not only in the order set (the number) but of the number type.

The solution set to the equation can be found from doubling the curve in (1) with an anti-holomorphic counterpart, finding the metric on the space which is dependent only on a and b , and generating the geodesic flows from a known solution such as $y^2 = b$ and $x = 0$ for b equaling a square.

The geodesic flow equations are represented by the two equations,

$$m = f(\tau; m_0, n_0) \quad n = g(\tau; m_0, n_0) , \quad (2)$$

together with their redundant conjugate equations. These equations require the metric on the elliptic curve in order to analyze, and the singularities label all of the disallowed integers m and n ; the latter represent non-allowed solutions to the curve. Systems of equations such as a hyperelliptic ones, or coupled equations in multi-variables can be analyzed in the same way.

The differential representation in (2) has a formal power series expansion,

$$m = \sum a_p(m_0, n_0) \tau^p \quad n = \sum b_q(m_0, n_0) \tau^q . \quad (3)$$

This series has to be analyzed, given the metric data, for the existence of a τ given a trivial solution such as $m_0 = n_0 = 0$. One condition is that one pair of the roots, pair meaning a root from both equations in (3), has to be real, non-zero, and identical. The coefficients $a_p(m_0, n_0)$ and $b_q(m_0, n_0)$ are derived from the metric data on the four-dimensional holomorphic and anti-holomorphic elliptic curves. In general, however,

the full functional dependence on τ in (3) should be analyzed including a possible analytic continuation (e.g. hypergeometric functions for example).

The group of integers are set into the polynomial equation

$$Q(x, m) = \prod_{n=0} (x - \sum h(m, n)) . \quad (4)$$

Another polynomial $P(x, m)$ can be defined which represents the complement set of integers, i.e. the allowed solution set to the elliptic equation,

$$P(x, m) = \frac{\prod_{m=0} (x - m)}{Q(x, m)} . \quad (5)$$

The polynomial in (5) generates all of the integer, or in general rational number solutions to the the curve,

$$y^2 = x^3 + ax + b . \quad (6)$$

The degree of the polynomial $P(x, m)$ in (5) generates the order of the solution set to (6).

Basically the functions

$$G_{m,n;m_0,n_0}^{(1)} = \sum a_p(m, n) \tau^p - m_0 \quad (7)$$

and

$$G_{m,n;m_0,n_0}^{(2)} = \sum b_q(m, n) \tau^q - n_0 . \quad (8)$$

or

$$G_{m,n;m_0,n_0} = G^{(1)} + iG^{(2)} , \quad (9)$$

are required for the number of zeros and singularities. These functions are found from the metric on the four-dimensional space spanned by the elliptic curve in (6) and its anti-holomorphic counterpart. These functions are generalized elliptic, or Appel functions, with a further generalization to non-elliptic coupled algebraic systems including hypergeometric ones. Standard differential or analysis techniques can find

the zeros and infinities, as well as their analytic continuations. The number of zeros of the functions, or of the complex one $G_{m,n;m_0,n_0}(\tau)$, are required to find the polynomial solutions to the elliptic curve in (1) and group them into the function in (5). The latter function seems clear to derive as well as its order.

The metric on these spaces is direct to find through a tree-level D-term analysis of the quiver formulation [4]. These metrics are found number theoretically through the pinching of classical ϕ^3 diagrams, through counting their number at zero momentum [5]. The geodesic flow solutions generating these functions are one-dimensional and are derivable [5]. General systems of polynomial equations are mapped to sets of functions in (8) depending on their number and dimensionality, i.e. variables in x and y . The analytic continuation, and their number of zeros and infinities, count the solutions in various number fields.

References

- [1] G. Chalmers, *Geometric Solutions to Algebraic Equations*, physics/0503175.
- [2] G. Chalmers, *A map from scalar field theory to integer polynomial solutions*, physics/0506013.
- [3] G. Chalmers, *Integer and Rational Solutions to Polynomial Equations*, physics/0503200.
- [4] G. Chalmers, *Tree amplitudes in scalar field theories*, physics/0504173.
- [5] G. Chalmers, *To appear*.